

Formulaire de trigonométrie

Les formules sont valables sous réserve qu'elles soient définies.

FONCTIONS CIRCULAIRES

$$\boxed{\cos^2 x + \sin^2 x = 1}$$

$$\cos x = \frac{e^{ix} + e^{-ix}}{2} \quad \sin x = \frac{e^{ix} - e^{-ix}}{2i}$$

$$e^{ix} = \cos x + i \sin x$$

$$\tan x = \frac{\sin x}{\cos x}$$

$$\cos(a+b) = \cos a \cos b - \sin a \sin b$$

$$\cos(a-b) = \cos a \cos b + \sin a \sin b$$

$$\sin(a+b) = \sin a \cos b + \cos a \sin b$$

$$\sin(a-b) = \sin a \cos b - \cos a \sin b$$

$$\tan(a+b) = \frac{\tan a + \tan b}{1 - \tan a \tan b}$$

$$\tan(a-b) = \frac{\tan a - \tan b}{1 + \tan a \tan b}$$

$$\cos a \cos b = \frac{1}{2} (\cos(a+b) + \cos(a-b))$$

$$\sin a \sin b = \frac{1}{2} (\cos(a-b) - \cos(a+b))$$

$$\sin a \cos b = \frac{1}{2} (\sin(a+b) + \sin(a-b))$$

$$\sin p + \sin q = 2 \sin \frac{p+q}{2} \cos \frac{p-q}{2}$$

$$\sin p - \sin q = 2 \cos \frac{p+q}{2} \sin \frac{p-q}{2}$$

$$\cos p + \cos q = 2 \cos \frac{p+q}{2} \cos \frac{p-q}{2}$$

$$\cos p - \cos q = -2 \sin \frac{p+q}{2} \sin \frac{p-q}{2}$$

$$\tan p + \tan q = \frac{\sin(p+q)}{\cos p \cos q}$$

$$\tan p - \tan q = \frac{\sin(p-q)}{\cos p \cos q}$$

$$\begin{aligned} \cos(2a) &= \cos^2 a - \sin^2 a = 2 \cos^2 a - 1 \\ &= 1 - 2 \sin^2 a \end{aligned}$$

$$\sin(2a) = 2 \sin a \cos a$$

$$\cos^2 a = \frac{1 + \cos(2a)}{2} \quad \sin^2 a = \frac{1 - \cos(2a)}{2}$$

$$\cos(2a) = \frac{1 - \tan^2 a}{1 + \tan^2 a} \quad \sin(2a) = \frac{2 \tan a}{1 + \tan^2 a}$$

$$\tan(2a) = \frac{2 \tan a}{1 - \tan^2 a}$$

$$\forall x \in [-1, 1], \operatorname{Arcsin} x + \operatorname{Arccos} x = \frac{\pi}{2}$$

$$\forall a \in \mathbb{R}^*, \operatorname{Arctan} a + \operatorname{Arctan} \frac{1}{a} = \operatorname{sgn} a \frac{\pi}{2}$$

$$\forall x \in]-1, 1[, \operatorname{Arcsin}' x = \frac{1}{\sqrt{1-x^2}}$$

$$\forall x \in]-1, 1[, \operatorname{Arccos}' x = \frac{-1}{\sqrt{1-x^2}}$$

$$\forall x \in \mathbb{R}, \operatorname{Arctan}' x = \frac{1}{1+x^2}$$

FONCTIONS HYPERBOLIQUES

$$\boxed{\operatorname{ch}^2 x - \operatorname{sh}^2 x = 1}$$

$$\operatorname{ch} x = \frac{e^x + e^{-x}}{2} \quad \operatorname{sh} x = \frac{e^x - e^{-x}}{2}$$

$$e^x = \operatorname{ch} x + \operatorname{sh} x$$

$$\operatorname{ch}(a+b) = \operatorname{ch} a \operatorname{ch} b + \operatorname{sh} a \operatorname{sh} b$$

$$\operatorname{ch}(a-b) = \operatorname{ch} a \operatorname{ch} b - \operatorname{sh} a \operatorname{sh} b$$

$$\operatorname{sh}(a+b) = \operatorname{sh} a \operatorname{ch} b + \operatorname{ch} a \operatorname{sh} b$$

$$\operatorname{sh}(a-b) = \operatorname{sh} a \operatorname{ch} b - \operatorname{ch} a \operatorname{sh} b$$

$$\operatorname{ch} a \operatorname{ch} b = \frac{1}{2} (\operatorname{ch}(a+b) + \operatorname{ch}(a-b))$$

$$\operatorname{sh} a \operatorname{sh} b = \frac{1}{2} (\operatorname{ch}(a+b) - \operatorname{ch}(a-b))$$

$$\operatorname{sh} a \operatorname{ch} b = \frac{1}{2} (\operatorname{sh}(a+b) + \operatorname{sh}(a-b))$$

$$\operatorname{sh} p + \operatorname{sh} q = 2 \operatorname{sh} \frac{p+q}{2} \operatorname{ch} \frac{p-q}{2}$$

$$\operatorname{sh} p - \operatorname{sh} q = 2 \operatorname{ch} \frac{p+q}{2} \operatorname{sh} \frac{p-q}{2}$$

$$\operatorname{ch} p + \operatorname{ch} q = 2 \operatorname{ch} \frac{p+q}{2} \operatorname{ch} \frac{p-q}{2}$$

$$\operatorname{ch} p - \operatorname{ch} q = 2 \operatorname{sh} \frac{p+q}{2} \operatorname{sh} \frac{p-q}{2}$$

$$\begin{aligned} \operatorname{ch}(2a) &= \operatorname{ch}^2 a + \operatorname{sh}^2 a = 2 \operatorname{ch}^2 a - 1 \\ &= 1 + 2 \operatorname{sh}^2 a \end{aligned}$$

$$\operatorname{sh}(2a) = 2 \operatorname{sh} a \operatorname{ch} a$$

$$\operatorname{ch}^2 a = \frac{1 + \operatorname{ch}(2a)}{2} \quad \operatorname{sh}^2 a = \frac{\operatorname{ch}(2a) - 1}{2}$$